

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,  
please do not report the images to the  
Problem Image Mailbox.

GENE 07413

# An efficient expression vector for stable expression in human liver cells\*

(HepG2; human polypeptide chain elongation factor 1 $\alpha$ ; Geneticin)Dong Wan Kim<sup>a</sup>, Takashi Harada<sup>b</sup>, Izumu Saito<sup>c</sup> and Tatsuo Miyamura<sup>b</sup>

<sup>a</sup>Department of Microbiology, Chang Won National University, Chang Won 641-773, Korea. Tel. (82-5) 5168-7462; <sup>b</sup>Department of Virology II, National Institute of Health, Toyama 1-23-1, Shinjuku-ku, Tokyo 162, Japan; and <sup>c</sup>Institute of Medical Sciences, University of Tokyo, Minato-ku, Tokyo 108, Japan. Tel. (81-3) 3443-8111

Received by Y. Sakaki: 25 March 1993; Revised/Accepted: 20 May/23 May 1993; Received at publishers: 16 July 1993

## SUMMARY

The expression of the bacterial neomycin resistance (Nm<sup>R</sup>)-encoding gene was examined under control of the promoter region of the gene encoding the human polypeptide chain elongation factor 1 $\alpha$  (EF-1 $\alpha$ ). In a human liver cell line, HepG2, the plasmid pEF321-*neo* directed higher expression of the bacterial *neo* gene than the Nm<sup>R</sup> gene expression vectors using other promoters, like the SV40 early region or thymidine kinase of Herpes simplex virus and SV40 early region promoter linked to human T-cell leukemia virus-1 enhancer sequences.

It is often difficult to establish human liver cell lines which support the permanent expression of transfected foreign genes. Many mammalian expression vectors which utilize various viral promoters and enhancers have been used in human liver cell lines. In many cases, however, their expression decreases after cell passages even if the high level expression is transiently observed. Especially, the efficiency of colony formation by human liver cells employing drug-resistance genes expressed by various expression vectors is extremely low when compared with other cell types.

Here, we describe a versatile expression vector, pEF321, useful for the establishment of human liver cell clones which permanently express desired genes. The pEF321 vector utilizes the promoter of *EF-1 $\alpha$*  gene (Uetsuki et al., 1989). We previously reported that this

potent promoter is highly active in almost all kinds of mammalian cells tested. They included not only fibroblasts but also lymphoid or neuronal cells (Kim et al., 1990). A plasmid pEF321-*neo* was constructed by inserting the bacterial Nm<sup>R</sup> gene under the EF- $\alpha$  promoter of pEF321 vector (Kim et al., 1990). In this study, we compared the efficiency of its colony formation with those of pSV2-*neo*, pTK-*neo* (or pKan2), pSR $\alpha$ -*neo* plasmids in human hepatoblastoma cell line, HepG2, in the presence of G418. The plasmids pSV2-*neo*, pTK-*neo* (or pKan2), pSR $\alpha$ -*neo* contain the simian virus (SV40) early region promoter, TK promoter of herpes simplex virus type 1 (Yates et al., 1984), SV40 early region promoter linked to human T cell leukemia virus-1 enhancer sequences (Takebe et al., 1988), respectively. As shown in Fig. 1 and Table I, the number of colonies formed by pEF321-*neo* was about 100-fold larger than those by pSV2-*neo* or pTK-*neo*, and about tenfold larger than pSR $\alpha$ -*neo*. The average size of colonies formed by pEF321-*neo* plasmid was also severalfold larger than that by other plasmids tested in this study. Our previous Northern blotting data suggested that larger cell colonies reflected higher expression of the transfected genes (data not shown). By the transfection of 1  $\mu$ g of pEF321-*neo* by Ca-phosphate co-precipitation method in HepG2 cells grown on a 6-cm

Correspondence to: Dr. T. Miyamura, Department of Virology II, National Institute of Health, Toyama 1-23-1, Shinjuku-ku, Tokyo 162, Japan. Tel. (81-3) 5285-1111; Fax (81-3) 5285-1161.

\*On request, the authors will supply detailed experimental evidence for conclusions reached in this Brief Note.

Abbreviations: EF-1 $\alpha$ , human polypeptide chain elongation factor 1 $\alpha$ ; EF-1 $\alpha$ , gene encoding EF-1 $\alpha$ ; G418, Geneticin; Nm, neomycin; <sup>R</sup>, resistance/resistant; SV40, simian virus 40; TK, thymidine kinase-encoding gene.




Fig. 1. Colony formation by HepG2 cells ( $1.6 \times 10^5$ ) transfected with 1 mg of plasmids pSV2-neo or pEF321-neo on 6 cm-plates. After 24 h,  $3 \times 10^5$  cells from each culture were transferred to 10 cm plates. After selection by 1.0 or 1.2 mg G418/ml in Dulbecco's modified Eagle's medium with 10% fetal bovine serum for four weeks, colonies were stained with 5% Giemsa solution.

TABLE I

Number of colonies<sup>a</sup> formed in HepG2 cells

Plasmid <sup>b</sup>	G418 (mg/ml)	
	0.8	1.0
pSV2-neo	0	0
pEF321-neo	42	8
pTK-neo	0	0
pSR $\alpha$ -neo	4	1

<sup>a</sup> $6 \times 10^4$  cells were replated in 6-cm plates and selected by 0.8 or 1.0 mg G418/ml in Dulbecco's modified Eagle's medium with 10% fetal bovine serum for 4 weeks. Colonies were counted after staining with 5% Giemsa solution (see Fig. 1).

<sup>b</sup>The HepG2 cells were transfected with each plasmid DNA (see Fig. 1).

plate, several hundred colonies were obtained. In fact, we could isolate many HepG2 cell clones which stably produced the hepatitis C virus (HCV) core protein using the pEF321-neo plasmid (data not shown). The stable expression of the HCV core protein was observed for at least 4 months. Although we did not examine its expression efficiency in the primary hepatocytes or the liver cells of other species, the pEF321 vector is thus useful for transient as well as stable expression of foreign genes in human liver cells.

#### ACKNOWLEDGEMENTS

We thank Y. Matsuura, T. Suzuki and A. Ando for helpful discussions. D.W.K. is a recipient of the Japanese government research award for foreign specialists from the Science and Technology Agency of the government of Japan.

#### REFERENCES

- Kim, D.W., Uetsuki, T., Kaziro, Y., Yamaguchi, N. and Sugano, S.: Use of the human elongation factor 1 $\alpha$  promoter as a versatile and efficient expression system. *Gene* 91 (1990) 217-223.
- Takebe, Y., Seiki, M., Fujisawa, J.-I., Hoy, P., Yokota, K., Arai, K.-I., Yoshida, M. and Arai, M.: SR $\alpha$  promoter: an efficient and versatile mammalian cDNA expression system composed of the simian virus 40 early promoter and the R-US segment of human T-cell leukemia virus type 1 long terminal repeat. *Mol. Cell. Biol.* 8 (1988) 466-472.
- Uetsuki, T., Naito, A., Nagata, S. and Kaziro, Y.: Isolation and characterization of the human chromosomal gene for polypeptide chain elongation factor 1 $\alpha$ . *J. Biol. Chem.* 264 (1989) 5791-5798.
- Yates, J., Warren, N., Reisman, D. and Sugden, B.: A cis-acting element from the Epstein-Barr viral genome that permits stable replication of recombinant plasmids in latently infected cells. *Proc. Natl. Acad. Sci. USA* 81 (1984) 3806-3810.